

Asian Journal of Research in Crop Science

Volume 8, Issue 4, Page 264-272, 2023; Article no.AJRCS.103281 ISSN: 2581-7167

Digital Technologies in Crop Genotype Designing Methods: Scope, Limitations and Future Perspectives

Aaron Chimbelya Siyunda ^{a,b*}, Emmanuel Chikalipa ^{a*}, Vinita Ramtekey ^c, Ntombokulunga Mbuma ^{d,e}, Mick Mwala ^a, Natasha Muchemwa Mwila ^a, Tesfaya Mitika Regassa ^f and Dyness Nshimbi ^b

 ^a Department of Plant Science, School of Agricultural Sciences, University of Zambia (UNZA), P.O. Box 32379, Great East Road Campus, Lusaka, Zambia.
 ^b Department of Crop Science, Natural Resources Development College (NRDC), P/Bag CH 99, Great East Road, Lusaka, Zambia.
 ^c ICAR-Indian Institute of Seed Science, Mau, Uttar Pradesh-275103, India.
 ^d Department of Plant Breeding, Agricultural Research Council, Vegetables, Industrial and Medicinal Plants Institute, Private Bag X293, Pretoria 0001, South Africa.
 ^e Department of Plant Sciences, Plant Breeding, University of the Free State, P.O. Box 339, Bloemfontein 9300, South Africa.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRCS/2023/v8i4206

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/103281

> Received: 22/05/2023 Accepted: 26/07/2023 Published: 19/08/2023

Review Article

ABSTRACT

The modern world agricultural sector has come under severe attack from several factors. These factors range from biotic to abiotic factors and they present threats to the environment and the world economies at large. If agricultural production is made more sustainable, it can be able to

*Corresponding author: E-mail: aaron.siyunda@yahoo.com, chikalipawesu@gmail.com;

Asian J. Res. Crop Sci., vol. 8, no. 4, pp. 264-272, 2023

combat the current food shortages. Looking into the present scenario, there is a great need to improve the traditional breeding designing methods to develop genotypes of different crops that would be able to withstand the current adverse effects brought about by persistent climate change. Central to the basis and key factor of improving the designing methods in crop production are different digital technologies such as Artificial Intelligence (AI), Deep Learning (DL), Machine Learning (ML), Geographical Information System (GIS), Precision Agriculture (PA), and Remote Sensing (RS). The digitalization of traditional breeding strategies has its weaknesses in terms of genetic gains it could offer in improving crop production. However, improving digital technologies would result in improved designing methods of crop production that would consequently result in increasing agricultural production and productivity. Therefore, the current review highlights the gains that have been made especially by AI and ML in designing methods of crop production. In addition, the review also highlights the limitations of these digital tools and their potential in crop designing methods for future crop genetic gains and production as well.


Keywords: Artificial intelligence; crop design methods; digitalization; machine learning; production.

1. INTRODUCTION

A re-look at how productivity and production in agriculture, are addressing the issue of food shortage currently and in the future is something that cannot be argued anymore. Especially that, the world's population is expected to rise by approximately 9.6 billion by 2050 [1]. Any efforts designed to overcome the current and future challenges that are affecting or have the potential to affect agricultural production and productivity negatively should be encouraged [2]. In today's agriculture sector, there are myriad factors that affect the full potential of crop production [3]. These factors range from biotic to abiotic stress conditions that have the potential to significantly affect crop development, from breeding to the production point. Major challenges experienced by agricultural production in the recent past include bottlenecks created by climate change, the reduction of water for irrigation purposes, an exponential rise in the cost of production, and the general decrease in the workforce dedicated to agricultural production as a result of the COVID-19 pandemic among other factors [4]. Both biotic and abiotic factors present threats to the environment and also to the world economies, concerning achieving sustainability of the present and future agricultural production and supply systems. To combat and overcome the factors that negatively affect crop production, there is a need to significantly innovate ways and means that will help in keeping pace with the current constant climate change [5-8]. The seemingly unavoidable question would be, how best do we produce enough and of good quality food meant for the ever-growing population estimated to reach approximately 9.6 billion by 2050?

Modern crop design methods have achieved tremendous gains at the breeding and also

production levels. Way back in the 19th century, genetic gains were heavily achieved through crossing and selection which involves combining and selecting classical breeding methods based on Mendel's laws, which also touches on experimental designs used in plant breeding [9]. During the 20th century, the gene pool enhancement that touches areas like wide crosses, pentaploid breeding in crops such as wheat, and also doubled haploid production were used to advance the breeding strategies. At the dawn of the 21st century, other crop designing methods came into the picture which includes rapid generation advancement e.g., shuttle breeding, and speed breeding (Fig 1), hybrid breeding, high throughput phenotyping platforms, high throughput genotyping platforms (genomics, pangenomics, transcriptomics, proteomics), epigenetics, and genome editing technologies [10]. Suffice to say, the advent of digital technologies is becoming a central point in designing methods that are promising to keep pace with the current constant and persistent climate change [11]. In the past, the development of crop designing methods was highly influenced by traditional or conventional methods. Currently, among all the digital technologies, AI and ML, have been employed on a larger scale to integrate with crop designing methods that are helping plant breeders to build genotypes that have high resilience towards climate change. Inspite of the gains made, the currently existing digital technologies have their weaknesses that if not worked on, may not be good enough for future research and crop sustainability production [12]. Therefore, this review becomes vital because it highlights the gains, limits, and potential that digital technologies (i.e., AI and ML) have in aiding crop designing methods (Fig 1) for future crop genotype designs, that could be able to overcome the current adverse



Siyunda et al.; Asian J. Res. Crop Sci., vol. 8, no. 4, pp. 264-272, 2023; Article no.AJRCS.103281

Fig. 1. Digital agriculture for crop genetic improvement

effects caused by climate change, to overcome the current food shortages.

2. ACHIEVEMENTS IN DESIGNING METHODS FOR FUTURE CROP GENOTYPES

2.1 Artificial Intelligence

Al has been defined by several researchers in different ways, central to its definition is the concept of integrating human intelligence in a machine, designed to operate in the same way humans do, by imitating their behavior [13]. Furthermore, researchers have highlighted that the term AI is mainly applied when describing a machine that can demonstrate the characteristics of human beings. It embraces the use of sensors and instruments such as spectrometers, digital cameras, etc. to enhance the process of capturing the data, before analysis and decisionmaking [14]. Previous strides of gains and achievements made by AI (Table 1) in the academic cycles have shown both opportunities and threats, strength and weakness, for various domains in science, including crop breeding programs. To achieve desired tasks effectively

and efficiently through proper machine Al methods require programming, large repositories of data. Besides, it uses algorithms that require large amounts of data to train the machine to enhance its ability in moderating the process of making decisions thereby promoting complex traits screening, such as detecting disease invasion in disease trial screening, at an early stage of crops and this, in turn, has improved the process of making decisions. In terms of developing crop designing methods, AI has made a significant contribution in the area of high-throughput phenotyping. This has been demonstrated by some researchers [5], which demonstrated the power of AI in giving insights into complex traits such as the architectural genetics of flowering time in wheat plants. Similarly, it has allowed and helped plant breeders to expand into areas that have not been explored in the past by designing crop methods that are capable of helping breeders to develop smart climate crop genotypes [3]. Previously, measuring phenotypic crop traits has been centered on traditional methods that involve destructive means succeeded by laboratory steps to explore phenotypic traits linking them to their genetic functions.

emergence of AI through The imaging techniques has recorded achievement in this area, in that, sampling of plants can be done in a non-destructive manner. capturing, by processing, and analyzing data for observable traits of interest in different crops [9]. Moreover, its application in imaging techniques, it has been widely used, in-field phenotyping platforms and open-source devices for the collection of data also in cyberinfrastructure for and data in different plant breeding management programs to come up with better crop designing methods that have high potential in facilitating the breeding of climate-smart elite lines [14]. Contrary to this, the use of AI has a to create high potential technological unemployment. AI-based models and tools used in different agricultural operations that are commercially available to ease the phenotyping, data collection, assessment of biotic and abiotic stress, and prediction of vield and quality of the products are shown in Tables 1 and 2.

2.2 Machine Learning

The focus of AI research has been ML since the 1970s. Different statistical models have since been developed to aid crop breeding programs, e.g., Bayesian networks, Perceptron, and support vector machines [27]. However, the challenge has been and still is that there is no single framework that works in a better way for all tasks. The determination of the best models for a given problem still poses a challenge. Despite the highlighted challenges in determining statistical models for a given task, ML models and algorithms still receive a wide and comprehensive utilization in plant breeding programs (Table 2). This may be due to the rising and successful establishment of data processing infrastructure, combined with the neutral network's establishment, high-level computational, and abundance of data [28]. There have been several gains and achievements made by machine learning in plant breeding programs (Table 1) [29]. Major achievements of machine learning models and algorithms in crop breeding include the exploration of large datasets and further discovering models, that are capable of developing resolutions through the development of patterns that combine features simultaneously unlike the process that analyses the features independently and separately. The step of combining the features simultaneously instead of

analyzing them separately used to be a challenge when using traditional processing methods, this may have been due to the high complexity of images of the plants and their huge quantity, which in this case, machine learning models can handle. ML has been effectively applied in the process of identification and classification of diseases that affect crops such as wheat, maize, and soybean, among other tasks [28]. To achieve this, supervised learning has been of help through the training of algorithms with images sampled from a large dataset, and in this way, algorithms can identify and classify plants that are diseased during disease screening [30]. The gains of ML regarding disease identification have been demonstrated in the research by Moshou et al.. 2004 [10]. The researchers successfully used ML to automatically detect yellow rust in wheat. Though ML has the strength to detect crop diseases as demonstrated, it has one central weakness, in that, it can prevent the exploration of unexpected and novel traits that less precise unsupervised machines can be able to discover in crops [31].

The establishment, enhancement, and advancement of high-throughput phenotyping that is nondestructive in crop breeding programs have heavily been influenced positively by imaging techniques of ML. Central to this gain is the ability of ML's utilization of multiple imaging sensors that can collect the data of plants in near real-time platforms [32]. These sensors have the capability of collecting huge amounts and volumes of phenotypic data that can easily be handled by ML, since it was built and has been developed to handle big data challenges, such as those experienced when dealing with traditional breeding methods in designing crop genotypes. Large volumes of data (big datasets), still pose a challenge with regard to being sorted out, analyzed, and interpreted when using traditional processing methods but ML models and algorithms have the power to utilize ways and approaches that are effective, faster, and efficient in the process of analyzing the data collected. Through its models and algorithms, has also made some gains in utilizing the tools such as statistics, decision and probability theories, and optimization in the process of extracting features from enormous datasets, thereby creating possibilities by making it easy to identify features in tasks that are complex such as phenotyping of different forms of stress in different crops [33].

Model	Application	Crop	Inputs	Results	Ref
DCNN	High Throughput Phenotyping	Wheat	Genotype and phenotype data	(Correlation Coefficient)>0.7	[15]
DCNN	High Throughput Phenotyping	Soybean	Genotype and phenotype data	(Correlation Coefficient)>0.4	[16]
DNN	Yield	Soybean	Unmanned Aerial System	R2.0.72	[17]
	Prediction		Images		
ANN	Yield	Wheat	Unmanned Aerial System	R2.0.7701, 0.112	[18]
	Prediction		based Vis		
RF	Yield	Wheat	Unmanned Aerial System	R2.0.7800, 0.1030	[18]
	Prediction		based Vis		
PLSR	Yield	Wheat	Unmanned Aerial System	R2.0.7667, 0.1353	[18]
	Prediction		based Vis		
DNN	Yield	Maize	2018 Syngenta Crop	Root Mean Square Error: 46%	[19]
	Prediction		Challenge	-	
RCNN	Fruit	Citrus	Unmanned Aerial System	Precision>90%	[20]
	detection				
RCNN	Fruit	Apples	Unmanned Aerial System	R2.0.80	[21]
	detection				
CNN	Weed	Rice	Unmanned Aerial System	General accuracy	[22]
	detection			weed mapping:89%	
				weed identifing:88%	
DCNN	Disease	Wheat	Unmanned Aerial System	Accuracy: 0.85	[23]
	Detection		images (hyperspectral)		
DCNN	Disease	Banana	Field images	General	[24]
	detection			Accuracy: >90%	
RF	Disease	Wheat	Unmanned Aerial System	Accuracy: 0.77	[23]
	detection		images (hyperspectral)		
RF	Biomass	Barley	Unmanned Aerial System	(Correlation Coefficient): 0.95	[25]
	Estimation		images		
CNN	Biomass	Maize	Field images	Accuracy: 99%	[26]
	Estimation				

Table 1. Artificial intelligence models used in different agricultural applications

DCNN; Deep Convolution Neutral Network, DNN; Deep Neutral Network, ANN; Artificial Neural Network, RF; Random Forest, PLSR; Partial Least Square Regression, RCNN; Region- Convolutional Neutral Network, CNN; Convolution Neural Network

Table 2. Artificial Intelligence	(AI)-Based tools commercial	v available for agriculture
	() =	,

Company	Products	Website
Trace Genomics	This product is a soil analysis system, it uses machine learning and is designed to provide a sense of the soil's strengths and weaknesses	tracegenomics.com
Harvest CROO Robotics	This product is a robot system designed for picking and packing vegetables.	harvestcroorobotics.com
Blue River Technology	The product is a smart farm machine, designed for the management of crops at a plant level and further protecting the crops from weed infestation.	bluerivertechnology.com
AGEYE Technologies	This product was specifically designed for indoor farming and it's an Artificial Intelligence-powered platform.	ageyetech.com
Fasal	This product uses affordable sensors, it's been designed to provide critical parameters to small farmers, central to this technology is Artificial Intelligence.	fasal.co
Vine View	This is the product that has aerial-based spectral sensors. It also has a cloud-based image processing service that is designed to monitor the plant health system. Its aerial-based spectral sensors are highly specialized.	vineview.com
PEAT	This is a product that is designed with a high specialization in image recognition i.e., identifying potential abnormalities and nutrient deficiencies in soils. Central to this product is deep learning.	plantix.net
HelioPas Al	This is a system developed to monitor soil moisture content and regulate the irrigation process. It is also used in the detection and management of mildew and drought.	heliopas.com
aWhere	This is the product that is designed for indoor farming. Central to this product is its Artificial Intelligence-powered platform.	awhere.com
FarmShort	This is the product designed to prescribe variables and carry out integrated scouting for farmers. Its core purpose is based on images captured using drones and satellites.	farmshots.com
Root Al	This is a product that is designed for indoor farmers. Central to this product is its Artificial Intelligence-powered platform that is automated and has robotic solutions.	root-ai.com
Ibex Automation	This is a product with a robust Agric-robotic system that is autonomous. It also has an autonomous system that can detect and spray weeds.	ibexautomation.co.uk

3. WAY FORWARD

With a lot of uncertainty that still covers today's environment due to the ever-changing climatic conditions [34], there is a need to optimize the utilization of digitalization, as this has a heavy bearing on the designs of breeding methods to be utilized for future plant breeding programs. The strength of artificial intelligence ranges from molecular to organismal scale, with this ability, combined with its capability to characterize complex traits in a very detailed and near realtime manner, AI still presents a ray of hope in designing crop methods that have the potential to bridge the gap between the phenotypic and genotypic traits of diverse crops.

Likewise, inspite of the gains and achievements made by crop designing methods using ML models and algorithms, there is a need to still explore ways of making it more effective and efficient to improve the development of climatesmart crop genotypes. Some of the strong points of ML are its ability to handle large volumes of data (big datasets), creating possibilities for developing and advancing ML approaches to a level where a single framework will be able to handle multiple tasks simultaneously and precisely which would otherwise enable future crop researchers to create crop designing methods that would have high accuracy, great precision and easy to execute. Furthermore, as feature identification of complex traits, it provides a platform to design crops with great precision and less time-consuming, and has the potential to reduce breeding cycles of crops to increase the genetic gains. Below is the chart describing the impact and potential of digital agriculture with a focus on AI and ML and their contribution in optimizing crop genetic improvement if efficiently and effectively used.

4. CONCLUSION

Innovations that can help us produce food sustainably, more effectively, and efficiently are no longer a question to be debated. Digital technologies such as AI and ML, present a promising platform for designing methods that have a high potential to make crop production operate at an optimal level where food production is concerned. Moving forward, there is a great need to improve the weaknesses that still exist in AI and ML if the current challenges and future needs are to be addressed. Improving digital technologies would result in an improvement in designing methods of crop production which would consequently increase agricultural production and productivity.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

 Siyunda AC, Chikalipa E, Mfune T, Hamvumba R. Digitalizing agriculture for sustainable crop production. IJSB. 2022; 11(1): 55-61. Available:https://doi.org/10.5281/zenodo.6

462310

Mwila NM, Munyinda K, Mwala M, Kamfwa K, Kambikambi T, Siyunda A, Sinyangwe S, Kanenga K, Alamu EO, Rubaihayo P. Situational analyses on cowpea value chain in Zambia: the case of an untapped legume, Cogent Food & Agriculture. 2022; 8(1):2094060,

Available:https://doi.org/10.1080/23311932 .2022.2094060

- Beans C. Crop researchers harness artificial intelligence to breed crops for the changing climate. Inner work; 2020. Available:www.pnas.org/cgi/doi/10.1073/p nas.2018732117
- 4. Furbank RT, Tester M. Phenomics-Technologies to relieve the phenotyping bottleneck. Trends Plant Sci. 2011;16: 635-644. Available:https://doi.org/10.1016/j.tplants.2

Available:https://doi.org/10.1016/j.tplants.2 011.09.005

5. Wang X, Xuan H, Evers B, Shrestha S, Pless R, Poland J. High-throughput phenotyping with deep learning gives insight into the genetic architecture of flowering time in wheat. Giga Sci. 2019; 8: 1-11.

Available:https://doi.org/10.1093/gigascien ce/giz120

- Siyunda AC, Mwila MN, Mwala M, Munyinda K, Kamfwa K, Chipabika G, Nshimbi D. Screening for resistance to cowpea aphids (*Aphis craccivora* Koch.) in mutation derived and cultivated cowpea (*Vigna unguiculata* L. Walp.) genotypes. Int J. Sci and Busi. 2022;13(1):15-26. Available:https://doi.org/10.5281/zenodo.6 614442
- Siyunda AC, Mwila MN, Mwala M, Munyinda K, Kamfwa K, Chipabika G, Nshimbi D. Laboratory screening of cowpea (*Vigna unguiculata* L. Walp.)

genotypes against pulse beetle (*Callosobruchus maculatus* F.). Int J. Sci. Busi. 2022;6(1):85-93. Available:https://doi.org/10.5281/zenodo.6 880440

- Siyunda AC, Mwila MN, Mwala M, Munyinda K, Kamfwa K, Chipabika G. Mutation Breeding: A modern tool for shelflife extension in Zambian cowpea lines. Journal of Genetics, Genomics & Plant Breeding. 2023;7(1):12-22. ISSN (Online): 2581-3293.
- Sabanci K, Toktas A, Kayabasi A. Grain classifier with computer vision using adaptive neuro-fuzzy inference system Pdf. J. Sci. Food Agric. 2017;97:3994-4000.

Available:https://doi.org/10.1002/jsfa.8264

 Moshou D, Bravo C, West C, Wahlen S, McCarteny A, Ramon H. Automated detection of "yellow rust" in wheat using reflectance measurements and neutral networks. Computer. Electron. Agric. 2004; 44:173-188.
 Available: https://doi.org/10.1016/i.compag

Available:https://doi.org/10.1016/j.compag. 2004.04.003

 Nabwire S, Suh HK, Kim MS, Baek I, Cho BK. Review: Application of Artificial intelligence in phenomics. Sensors. 2021; 21;4363. Available:https://doi.org/10.3390/s2113436

Available:https://doi.org/10.3390/s2113436

 Sabanci K, Kayabasi A, Toktas A. Computer vision-based method for classification of wheat grains using artificial neural network. J. Sci. Food Agric. 2017; 97: 2588-2593.

Available:https://doi.org/10.1002/jsfa.8080

- Jensen T, Apan A, Young E, Zeller L. Detecting the attributes of a wheat Crop using digital imagery acquired from a lowaltitude platform. Computer. Electron. Agric. 2007;59:66-77. Available:https://doi.org/10.1016/j.compag. 2007.05.004
- Munns R, James RA, Smault XRR, Furbank RT, Jones HG. New phenotyping methods for screening wheat and barley for beneficial responses to water deficit. J. exp. Bot. 2010;61:3499-3507. Available:https://doi.org/10.1093/jxb/erq19 9
- Ma W, Qiu Z, Song J, Li J, Cheng Q, Zhi J, Ma C. A deep convolutional neural network approach for predicting phenotypes from genotypes. Planta. 2018;248:1307-1318.

Available:https://doi.org/10.1007/s00425-018-2976-9

- Liu Y, Wang D, He F, Wang J, Joshi T, Xu D. Phenotype prediction and genome-wide association study using deep convolutional neural network of soybean. Front Genet. 2019;10:1091 Available:https://doi.org/10.3389/fgene.201
- 9.01091
 17. Maimaitijiang M, Sagan V, Sidike P, Hartling S, Esposito F, Fritschi FB. Soyabean yield prediction from UAV using multimodal data fusion and deep learning. Remote Sens Environ. 2020;237;111599. Available:https://doi.org/10.1016/j.rse.2019 .111599
- Fu Z, Jiang J, Gao Y, Krienke B, Wang M, Zhong K, Cao Q, Tian Y, Zhu Y, Cao W, Liu X. Wheat growth monitoring and yield estimation based on multi-rotor unmanned aerial vehicle. Remote Sens. 2020;12; 508. Available:https://doi.org/10.3390/rs120305

Available:https://doi.org/10.3390/rs120305 08

- Khaki S, Wang L. Crop yield prediction using deep neural networks. Front Plant Sci. 2019;10:621. Available:https://doi.org/10.3389/fpls.2019. 00621
- Apolo-Apolo OE, Martinez-Guanter J, Egea G, Raja P, Perez-Ruiz M. Deep Learning Techniques for Estimation of the yield and size of the citrus fruits using UAV. Eur J Agron. 2020;115:126030. Available:https://doi.org/10.1016/j.eja.2020 .126030
- 21. Apolo-Apolo OE, Perez-Ruiz M, Martinez-Guanter J, Valente J. A cloud-based environment for generating yield estimation maps from apple orchards using UAV imagery and deep learning techniques. Front Plant Sci. 2020; 11:1086. Available:https://doi.org/10.3389/fpls.2020.

01086. Huang H, Deng J, Lan Y, Yang A, Deng X,

- Huang H, Deng J, Lan Y, Yang A, Deng X, Zang L. A fully convolutional network for weed mapping of unmanned aerial vehicle (UAV) imagery. PLoS One. 2018;13: e0196302. Available:https://doi.org/10.1371/journal.po ne.0196302
- 23. Zhang X, Han L, Dong Y, Shi Y, Huang W, Han L, Gonzalez-Moreno P, Ma H, Ye H, Sobeih TA. Deep learning-based approach for automated yellow rust disease detection from high-resolution

hyperspectral UAV images. Remote Sensi. 2019;11:1554. Available:https://doi.org/10.3390/rs111315

Available:https://doi.org/10.3390/rs111315 54

- Selvaraj MG, Vergaara A, Ruiz H, Safari N, Elayabalan S, Ocimati W, Biomme G. Al-powered banana diseases and pest detection. Plant Meth. 2019;15:92. Available:https://doi.org/10.1186/s13007-019-0475-z
- Nasi R, Viijanen N, Kairosoja J, Alhonoja K, Hakala T, Markelin L, Honkavaara E. Estimating biomass and nitrogen amount of barley and grass using UAV and aircraft based spectral and photogrammatic 3D features. Remote Sens. 2018;10:1082. Available:https://doi.org/10.3390/rs100710 82
- 26. Wiesner-Hanks T, Wu H, Stewart E, DeChat Kaczmar N, Lipson H, Gore MA, Nelson RJ. Millimeter-level plant diseases detection from aerial photographs via deep learning and crowd sourced data. Front Plant Sci. 2019;10:1082. Available:https://www.frontiersin.org/article
- s/10.3389/fpls.2019.01550/full
 27. Gianola D, Okut H, Weigel KA, Rosa GJ. Predicting complex quantitative traits with Bayesian neural networks: A case study with Jersey cows and wheat. BMC Genet. 2011;12:87.

Available:https://doi.org/10.1186/1471-2156-12-87

 Espositos S, Carputo D, Cardi T, Tripodi P. Applications and trends of machine learning in genomics and phenomics for next-generation breeding. Plants. 2020;9 34:

Available:https://doi:10.3390/plants901003 4

29. Jung J, Maeda M, Chang A, Bhandari M, Ashapure A, Landivar-Bowles J. The potential of remote sensing and artificial intelligence as tools to improve the resilience of agriculture production systems. Curr. Opinions in Biotech. 2021; 70:15-22.

Available:https://doi.org/10.1016/j.copbio.2 020.09.003

 Perez-Sanz F, Navarro PJ, Egea-Cortines M. Plant phenomics: An overview of image acquisition technologies and image data analysis algorithms. Giga Sci. 2017;6: 1-18.

Available:http://doi.org/10.1093/gigascienc e/gix092

- Sadeghi-Tehran P, Sabermanesh K, Virlet N, Hawkesford MJ. Automated method to determine two critical growth stages off wheat: Heading and flowering. Front. Plant Sci. 2017;8:1-14. Available:https://doi.org/10.3389/fpls.2017.00252
- Prey L, Von Bloh M, Schmidhalter U. Evaluating RGB imaging and multispectral active and hyperspectral passive sensing for assessing early plant vigour in winter wheat. Sens. 2018.18:2931. Available:https://doi.org/10.3390/s1809293 1
- Vigneau N, Ecarnot M, Rabatel G, Raumet P. Potential of field hyperspectral imaging as a non-destructive method to assess leaf nitrogen content in wheat. Field Crops Res. 2011:122;25-31. Available:https://doi.org/10.1016/j.fcr.2011. 02.003
- Mwila MN, Mukonze B, Siyunda A, Munyinda A. Climate Smart Crop: Evaluation of Selected Mutant Cowpea Genotypes for Yield, Earliness and Ground Cover in Eastern Zambia. Medicon Agriculture & Environmental Sciences. 2022;3(3):33-41.

© 2023 Siyunda et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/103281